



## **Passive dosing as a tool to derive fugacity capacities of a variety of leaves for semi-volatile persistent organic contaminants**

**Bolinus, D. J.; Mayer, Philipp; MacLeod, M.; Mclachlan, Michael S.; Jahnke, A.**

*Published in:*  
Science Across Bridges, Borders and Boundaries

*Publication date:*  
2014

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Bolinus, D. J., Mayer, P., MacLeod, M., Mclachlan, M. S., & Jahnke, A. (2014). Passive dosing as a tool to derive fugacity capacities of a variety of leaves for semi-volatile persistent organic contaminants. In *Science Across Bridges, Borders and Boundaries: Programme Book SETAC Europe*.

---

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**TH071 Passive dosing as a tool to derive fugacity capacities of a variety of leaves for semi-volatile persistent organic contaminants.**

D.J. Bolinius, Department of applied environmental science; P. Mayer, Technical University of Denmark / Department of Environmental Engineering; M. MacLeod, ITM Stockholm University / Dept of Applied Environmental Science ITM; M.S. McLachlan, Stockholm University; A. Jahnke, Stockholm University / Department of Applied Environmental Science ITM. The silicone polydimethylsiloxane (PDMS) has proven useful as a tool in experimental environmental chemistry: It has been applied successfully as a passive sampler<sup>1</sup> as well as a reliable donor in passive dosing experiments<sup>2</sup>. In this study we extend the application of PDMS as a passive dosing source to derive fugacity capacities ( $Z$ ) of a variety of matrices for semi-volatile persistent organic contaminants.  $Z$ , expressed in  $\text{mol m}^{-3} \text{Pa}^{-1}$ , is a measure of the extent to which a chemical can be absorbed by a given matrix at a specific temperature. It is an important and frequently used parameter in modeling the environmental fate of chemicals<sup>3</sup>. Our experimental apparatus, based on Mayer et al.<sup>4</sup>, consists of a thin disc of the sample wedged between 2 layers of PDMS loaded with the chemicals of interest. The PDMS is left to equilibrate with the sample and based on the known  $Z$  values of the PDMS for these chemicals, the  $Z$  value of the sample can be derived. The ease of the setup makes the approach suitable to measure the  $Z$  values of a wide variety of matrices. The challenge lies in keeping the PDMS the dominant reservoir of the chemical in the system. Analyzing tissues that are too rich in lipid risks depleting the PDMS to the extent that it becomes hard to accurately determine the fugacity capacity. Within the scope of our studies focusing on the so-called 'forest filter effect' we have applied the method to a variety of leaves. Existing multimedia chemical fate models that incorporate a forest compartment currently only assign two different  $Z$  values for foliage: One for conifers one for deciduous trees<sup>5</sup>. The composition of leaves is highly variable however. By measuring  $Z$  for a wide variety of foliage and linking these values to species-specific properties, we hope to achieve a better understanding of the cycling of semi-volatile persistent organic chemicals in forest systems. New  $Z$  values can easily be incorporated into existing fugacity based models or translated to leaf/air partitioning ratios. **References:** Reichenberg, F. et al. (2008). *Chem. Cent. J.* **2**, 8 Smith, K. E. C. et al. (2010). *Chem. Res. Toxicol.* **23**, 55–65. Mackay, D. (2001). *Multimedia Environmental Models: The Fugacity Approach*. Lewis Publishers, Chelsea, MI. Mayer, P. et al. (2005). *Environ. Sci. Technol.* **39**, 6123–6129. Wania, F. et al. (2006). *Environ. Modell. Softw.* **21**, 868–884.